

Different Approaches to Morse-Bott Homology

David Hurtubise
with Augustin Banyaga

Penn State Altoona
math.aa.psu.edu

Université Cheikh Anta Diop de Dakar, Senegal
May 19, 2012

Computing homology using critical points and flow lines

Perturbations

- Generic perturbations

- Applications of the perturbation approach

- A more explicit perturbation

Spectral sequences

- Filtrations and Morse-Bott functions

- Filtrations and spectral sequences

- Applications of the spectral sequence approach

Cascades

- Picture of a 3-cascade

- Applications of the cascade approach

- Cascades and perturbations

Multicomplexes

- Definition and assembly

- Multicomplexes and spectral sequences

- The Morse-Bott-Smale multicomplex

The Morse-Smale-Witten chain complex

Let $f : M \rightarrow \mathbb{R}$ be a Morse-Smale function on a compact smooth Riemannian manifold M of dimension $m < \infty$, and assume that orientations for the unstable manifolds of f have been chosen. Let $C_k(f)$ be the free abelian group generated by the critical points of index k , and let


$$C_*(f) = \bigoplus_{k=0}^m C_k(f).$$

Define a homomorphism $\partial_k : C_k(f) \rightarrow C_{k-1}(f)$ by

$$\partial_k(q) = \sum_{p \in \text{Cr}_{k-1}(f)} n(q, p)p$$


where $n(q, p)$ is the number of gradient flow lines from q to p counted with sign. The pair $(C_*(f), \partial_*)$ is called the **Morse-Smale-Witten chain complex** of f .

The height function on the 2-sphere



$$\begin{array}{ccccccc}
 C_2(f) & \xrightarrow{\partial_2} & C_1(f) & \xrightarrow{\partial_1} & C_0(f) & \longrightarrow & 0 \\
 \updownarrow \approx & & \updownarrow \approx & & \updownarrow \approx & & \\
 \langle n \rangle & \xrightarrow{\partial_2} & \langle 0 \rangle & \xrightarrow{\partial_1} & \langle s \rangle & \longrightarrow & 0
 \end{array}$$

The height function on a deformed 2-sphere




$$\begin{array}{ccccccc}
 C_2(f) & \xrightarrow{\partial_2} & C_1(f) & \xrightarrow{\partial_1} & C_0(f) & \longrightarrow & 0 \\
 \uparrow \approx & & \uparrow \approx & & \uparrow \approx & & \\
 \langle r, s \rangle & \xrightarrow{\partial_2} & \langle q \rangle & \xrightarrow{\partial_1} & \langle p \rangle & \longrightarrow & 0
 \end{array}$$


References for Morse homology

- ▶ Augustin Banyaga and David Hurtubise, **Lectures on Morse homology**, Kluwer Texts in the Mathematical Sciences **29**, Kluwer Academic Publishers Group, 2004.
- ▶ Andreas Floer, *Witten's complex and infinite-dimensional Morse theory*, J. Differential Geom. **30** (1989), no. 1, 207–221.
- ▶ John Milnor, **Lectures on the h-cobordism theorem**, Princeton University Press, 1965.
- ▶ Matthias Schwarz, **Morse homology**, Progress in Mathematics **111**, Birkhäuser, 1993.
- ▶ Edward Witten, *Supersymmetry and Morse theory*, J. Differential Geom. **17** (1982), no. 4, 661–692.

A Morse-Bott function on the 2-sphere



A Morse-Bott function on the 2-sphere



Can we construct a chain complex for this function? a spectral sequence? a multicomplex?

Generic perturbations

Theorem (Morse 1932)

Let M be a finite dimensional smooth manifold. Given any smooth function $f : M \rightarrow \mathbb{R}$ and any $\varepsilon > 0$, there is a Morse function $g : M \rightarrow \mathbb{R}$ such that $\sup\{|f(x) - g(x)| \mid x \in M\} < \varepsilon$.

Generic perturbations

Theorem (Morse 1932)

Let M be a finite dimensional smooth manifold. Given any smooth function $f : M \rightarrow \mathbb{R}$ and any $\varepsilon > 0$, there is a Morse function $g : M \rightarrow \mathbb{R}$ such that $\sup\{|f(x) - g(x)| \mid x \in M\} < \varepsilon$.

Theorem

Let M be a finite dimensional compact smooth manifold. The space of all C^r Morse functions on M is an open dense subspace of $C^r(M, \mathbb{R})$ for any $2 \leq r \leq \infty$ where $C^r(M, \mathbb{R})$ denotes the space of all C^r functions on M with the C^r topology.

Generic perturbations

Theorem (Morse 1932)

Let M be a finite dimensional smooth manifold. Given any smooth function $f : M \rightarrow \mathbb{R}$ and any $\varepsilon > 0$, there is a Morse function $g : M \rightarrow \mathbb{R}$ such that $\sup\{|f(x) - g(x)| \mid x \in M\} < \varepsilon$.

Theorem

Let M be a finite dimensional compact smooth manifold. The space of all C^r Morse functions on M is an open dense subspace of $C^r(M, \mathbb{R})$ for any $2 \leq r \leq \infty$ where $C^r(M, \mathbb{R})$ denotes the space of all C^r functions on M with the C^r topology.

Why not just perturb the Morse-Bott function $f : M \rightarrow \mathbb{R}$ to a Morse function?

The Chern-Simons functional

Let $P \rightarrow N$ be a (trivial) principal $SU(2)$ -bundle over an oriented closed 3-manifold N , and let \mathcal{A} be the space of connections on P . Define $CS : \mathcal{A} \rightarrow \mathbb{R}$ by

$$CS(A) = \frac{1}{4\pi^2} \int_M \text{tr} \left(\frac{1}{2} A \wedge dA + \frac{1}{3} A \wedge A \wedge A \right).$$

The above functional descends to a function $cs : \mathcal{A}/\mathcal{G} \rightarrow \mathbb{R}/\mathbb{Z}$ whose critical points are gauge equivalence classes of flat connections. Extending everything to $P \times \mathbb{R} \rightarrow N \times \mathbb{R}$, the gradient flow equation becomes the instanton equation

$$F + *F = 0,$$

where F denotes the curvature and $*$ is the Hodge star operator.

Instanton homology

Andreas Floer, *An instanton-invariant for 3-manifolds*, Comm. Math. Phys. **118** (1988), no. 2, 215–240.

Theorem. When N is a homology 3-sphere the Chern-Simons functional can be perturbed so that it has discrete critical points and defines \mathbb{Z}_8 -graded homology groups $I_*(N)$ analogous to the Morse homology groups.

Instanton homology

Andreas Floer, *An instanton-invariant for 3-manifolds*, Comm. Math. Phys. **118** (1988), no. 2, 215–240.

Theorem. When N is a homology 3-sphere the Chern-Simons functional can be perturbed so that it has discrete critical points and defines \mathbb{Z}_8 -graded homology groups $I_*(N)$ analogous to the Morse homology groups.

Generalizations: Donaldson polynomials for 4-manifolds with boundary, knot homology groups

The symplectic action functional

Let (M, ω) be a closed symplectic manifold and $S^1 = \mathbb{R}/\mathbb{Z}$. A time-dependent Hamiltonian $H : M \times S^1 \rightarrow \mathbb{R}$ determines a time-dependent vector field X_H by

$$\omega(X_H(x, t), v) = v(H)(x, t) \text{ for } v \in T_x M.$$

Let $\mathcal{L}(M)$ be the space of free contractible loops on M and

$$\tilde{\mathcal{L}}(M) = \{(x, u) \mid x \in \mathcal{L}(M), u : D^2 \rightarrow M \text{ such that } u(e^{2\pi i t}) = x(t)\} / \sim$$

its universal cover with covering group $\pi_2(M)$. The symplectic action functional $a_H : \tilde{\mathcal{L}}(M) \rightarrow \mathbb{R}$ is defined by

$$a_H((x, u)) = \int_{D^2} u^* \omega + \int_0^1 H(x(t), t) dt.$$

The Arnold conjecture

Andreas Floer, *Symplectic fixed points and holomorphic spheres*,
Comm. Math. Phys. **120** (1989), no. 4, 575–611.

Theorem. Let (P, ω) be a compact symplectic manifold. If I_ω and I_c are proportional, then the fixed point set of every exact diffeomorphism of (P, ω) satisfies the Morse inequalities with respect to any coefficient ring **whenever it is nondegenerate**.

The Arnold conjecture

Andreas Floer, *Symplectic fixed points and holomorphic spheres*,
 Comm. Math. Phys. **120** (1989), no. 4, 575–611.

Theorem. Let (P, ω) be a compact symplectic manifold. If I_ω and I_c are proportional, then the fixed point set of every exact diffeomorphism of (P, ω) satisfies the Morse inequalities with respect to any coefficient ring **whenever it is nondegenerate**.

Generalizations: Allowing H to be degenerate (e.g. $H = 0$) leads to critical submanifolds and Morse-Bott homology.

Lagrangian intersection homology

Let $L \subset P$ be a Lagrangian submanifold of a symplectic manifold (P, ω) and $\phi_1 : P \rightarrow P$ a Hamiltonian diffeomorphism such that $\phi_1(L)$ intersects L transversally. There is a Floer chain complex with chain groups generated by the elements of $L \cap \phi_1(L)$ and whose boundary operator is given by counting J -holomorphic curves $\mathbb{C}P^1 \rightarrow P$ from L to $\phi_1(L)$.

Lagrangian intersection homology

Let $L \subset P$ be a Lagrangian submanifold of a symplectic manifold (P, ω) and $\phi_1 : P \rightarrow P$ a Hamiltonian diffeomorphism such that $\phi_1(L)$ intersects L transversally. There is a Floer chain complex with chain groups generated by the elements of $L \cap \phi_1(L)$ and whose boundary operator is given by counting J -holomorphic curves $\mathbb{C}P^1 \rightarrow P$ from L to $\phi_1(L)$.

Theorem. (Floer 1988) If P is a compact symplectic manifold with $\pi_2(P) = 0$ and ϕ is an exact diffeomorphism of P **all of whose fixed points are nondegenerate**, then the number of fixed points of ϕ is greater than or equal to the sum of the \mathbb{Z}_2 -Betti numbers of P .

Lagrangian intersection homology

Let $L \subset P$ be a Lagrangian submanifold of a symplectic manifold (P, ω) and $\phi_1 : P \rightarrow P$ a Hamiltonian diffeomorphism such that $\phi_1(L)$ intersects L transversally. There is a Floer chain complex with chain groups generated by the elements of $L \cap \phi_1(L)$ and whose boundary operator is given by counting J -holomorphic curves $\mathbb{C}P^1 \rightarrow P$ from L to $\phi_1(L)$.

Theorem. (Floer 1988) If P is a compact symplectic manifold with $\pi_2(P) = 0$ and ϕ is an exact diffeomorphism of P **all of whose fixed points are nondegenerate**, then the number of fixed points of ϕ is greater than or equal to the sum of the \mathbb{Z}_2 -Betti numbers of P .

Generalizations: Extensive work by Fukaya, Oh, Ohta, and Ono to include the Morse-Bott case using **spectral sequences** and by Frauenfelder using **cascades**.

An explicit perturbation of $f : M \rightarrow \mathbb{R}$

Let T_j be a small tubular neighborhood around each connected component $C_j \subseteq Cr(f)$ for all $j = 1, \dots, l$. Pick a positive Morse function $f_j : C_j \rightarrow \mathbb{R}$ and extend f_j to a function on T_j by making f_j constant in the direction normal to C_j for all $j = 1, \dots, l$.

An explicit perturbation of $f : M \rightarrow \mathbb{R}$

Let T_j be a small tubular neighborhood around each connected component $C_j \subseteq Cr(f)$ for all $j = 1, \dots, l$. Pick a positive Morse function $f_j : C_j \rightarrow \mathbb{R}$ and extend f_j to a function on T_j by making f_j constant in the direction normal to C_j for all $j = 1, \dots, l$.

Let $\tilde{T}_j \subset T_j$ be a smaller tubular neighborhood of C_j with the same coordinates as T_j , and let ρ_j be a smooth bump function which is constant in the coordinates parallel to C_j , equal to 1 on \tilde{T}_j , equal to 0 outside of T_j , and decreases on $T_j - \tilde{T}_j$ as the coordinates move away from C_j .

An explicit perturbation of $f : M \rightarrow \mathbb{R}$

Let T_j be a small tubular neighborhood around each connected component $C_j \subseteq Cr(f)$ for all $j = 1, \dots, l$. Pick a positive Morse function $f_j : C_j \rightarrow \mathbb{R}$ and extend f_j to a function on T_j by making f_j constant in the direction normal to C_j for all $j = 1, \dots, l$.

Let $\tilde{T}_j \subset T_j$ be a smaller tubular neighborhood of C_j with the same coordinates as T_j , and let ρ_j be a smooth bump function which is constant in the coordinates parallel to C_j , equal to 1 on \tilde{T}_j , equal to 0 outside of T_j , and decreases on $T_j - \tilde{T}_j$ as the coordinates move away from C_j . For small $\varepsilon > 0$ (and a careful choice of the metric) this determines a Morse-Smale function

$$h_\varepsilon = f + \varepsilon \left(\sum_{j=1}^l \rho_j f_j \right).$$

Critical points of the perturbed function

If $p \in C_j$ is a critical point of $f_j : C_j \rightarrow \mathbb{R}$ of index λ_p^j , then p is a critical point of h_ε of index

$$\lambda_p^{h_\varepsilon} = \lambda_j + \lambda_p^j$$

where λ_j is the Morse-Bott index of C_j .

Critical points of the perturbed function

If $p \in C_j$ is a critical point of $f_j : C_j \rightarrow \mathbb{R}$ of index λ_p^j , then p is a critical point of h_ε of index

$$\lambda_p^{h_\varepsilon} = \lambda_j + \lambda_p^j$$

where λ_j is the Morse-Bott index of C_j .

Theorem (Morse-Bott Inequalities)

Let $f : M \rightarrow \mathbb{R}$ be a Morse-Bott function on a finite dimensional oriented compact smooth manifold, and assume that all the critical submanifolds of f are orientable. Then there exists a polynomial $R(t)$ with non-negative integer coefficients such that

$$MB_t(f) = P_t(M) + (1 + t)R(t).$$

Critical points of the perturbed function

If $p \in C_j$ is a critical point of $f_j : C_j \rightarrow \mathbb{R}$ of index λ_p^j , then p is a critical point of h_ε of index

$$\lambda_p^{h_\varepsilon} = \lambda_j + \lambda_p^j$$

where λ_j is the Morse-Bott index of C_j .

Theorem (Morse-Bott Inequalities)

Let $f : M \rightarrow \mathbb{R}$ be a Morse-Bott function on a finite dimensional oriented compact smooth manifold, and assume that all the critical submanifolds of f are orientable. Then there exists a polynomial $R(t)$ with non-negative integer coefficients such that

$$MB_t(f) = P_t(M) + (1 + t)R(t).$$

(Different orientation assumptions in [Banyaga-H 2009] than the proof using the Thom Isomorphism Theorem.)

The main idea behind the Banyaga-H proof

$$\begin{aligned}
 MB_t(f) &= \sum_{j=1}^l P_t(C_j)t^{\lambda_j} \\
 &= \sum_{j=1}^l \left(M_t(f_j) - (1+t)R_j(t) \right) t^{\lambda_j} \\
 &= \sum_{j=1}^l M_t(f_j)t^{\lambda_j} - (1+t) \sum_{j=1}^l R_j(t)t^{\lambda_j} \\
 &= M_t(h) - (1+t) \sum_{j=1}^l R_j(t)t^{\lambda_j} \\
 &= P_t(M) + (1+t)R_h(t) - (1+t) \sum_{j=1}^l R_j(t)t^{\lambda_j}
 \end{aligned}$$

The filtration associated to a Morse-Bott function

For a Morse-Bott function $f : M \rightarrow \mathbb{R}$ and $t \in \mathbb{R}$ we can define the “half-space” $M^t = \{x \in M \mid f(x) \leq t\}$.

If the critical values of f are $c_1 < c_2 < \dots < c_k$, then we have a filtration

$$\emptyset \subseteq M^{c_1} \subseteq M^{c_2} \subseteq \dots \subseteq M^{c_k}.$$

For any $j = 2, \dots, k$, M^{c_j} is homotopic to $M^{c_{j-1}}$ with a λ -disk bundle attached for each critical submanifold of index λ in the critical level $f^{-1}(c_j)$.

This is a generalization of the fact that a Morse function on M determines a CW-complex X that is homotopic to M .

The filtration associated to a Morse-Bott function

For a Morse-Bott function $f : M \rightarrow \mathbb{R}$ and $t \in \mathbb{R}$ we can define the “half-space” $M^t = \{x \in M \mid f(x) \leq t\}$.

If the critical values of f are $c_1 < c_2 < \dots < c_k$, then we have a filtration

$$\emptyset \subseteq M^{c_1} \subseteq M^{c_2} \subseteq \dots \subseteq M^{c_k}.$$

For any $j = 2, \dots, k$, M^{c_j} is homotopic to $M^{c_{j-1}}$ with a λ -disk bundle attached for each critical submanifold of index λ in the critical level $f^{-1}(c_j)$.

This is a generalization of the fact that a Morse function on M determines a CW-complex X that is homotopic to M .

The Morse-Smale-Witten boundary operator is defined differently than the boundary operator induced by the connecting homomorphism in the long exact sequence of a triple.

The spectral sequence associated to a filtration

Let (C_*, ∂) a filtered chain complex that is bounded below by $s = 0$. That is, suppose that we have a filtration

$$F_0 C_* \subset \cdots \subset F_{s-1} C_* \subset F_s C_* \subset F_{s+1} C_* \subset \cdots$$

where $F_s C_*$ is a chain subcomplex of C_* for all s . Define

$$\begin{aligned} Z_{s,t}^r &= \{c \in F_s C_{s+t} \mid \partial c \in F_{s-r} C_{s+t-1}\} \\ Z_{s,t}^\infty &= \{c \in F_s C_{s+t} \mid \partial c = 0\}. \end{aligned}$$

The bigraded R -modules in the spectral sequence associated to the filtration are defined to be

$$\begin{aligned} E_{s,t}^r &= Z_{s,t}^r / \left(Z_{s-1,t+1}^{r-1} + \partial Z_{s+r-1,t-r+2}^{r-1} \right) \\ E_{s,t}^\infty &= Z_{s,t}^\infty / \left(Z_{s-1,t+1}^\infty + (\partial C_{s+t+1} \cap F_s C_{s+t}) \right). \end{aligned}$$

The differentials in the spectral sequence

The differential $d^r : E_{s,t}^r \rightarrow E_{s-r,t+r-1}^r$ in the spectral sequence associated to a filtered chain complex is defined by the following diagram.

$$\begin{array}{ccc}
 Z_{s,t}^r & \xrightarrow{\quad \partial \quad} & Z_{s-r,t+r-1}^r \\
 \downarrow & & \downarrow \\
 Z_{s,t}^r / (Z_{s-1,t+1}^{r-1} + \partial Z_{s+r-1,t-r+2}^{r-1}) & \xrightarrow{\quad d^r \quad} & Z_{s-r,t+r-1}^r / (Z_{s-r-1,t+r}^{r-1} + \partial Z_{s-1,t+1}^{r-1})
 \end{array}$$

The R -module $E_{s,t}^r$ is isomorphic to $\bar{Z}_{s,t}^{r-1} / \bar{B}_{s,t}^{r-1}$ via an isomorphism given by the Noether Isomorphism Theorem.

The differentials in the spectral sequence

The differential $d^r : E_{s,t}^r \rightarrow E_{s-r,t+r-1}^r$ in the spectral sequence associated to a filtered chain complex is defined by the following diagram.

$$\begin{array}{ccc}
 Z_{s,t}^r & \xrightarrow{\quad \partial \quad} & Z_{s-r,t+r-1}^r \\
 \downarrow & & \downarrow \\
 Z_{s,t}^r / (Z_{s-1,t+1}^{r-1} + \partial Z_{s+r-1,t-r+2}^{r-1}) & \xrightarrow{\quad d^r \quad} & Z_{s-r,t+r-1}^r / (Z_{s-r-1,t+r}^{r-1} + \partial Z_{s-1,t+1}^{r-1})
 \end{array}$$

The R -module $E_{s,t}^r$ is isomorphic to $\bar{Z}_{s,t}^{r-1} / \bar{B}_{s,t}^{r-1}$ via an isomorphism given by the Noether Isomorphism Theorem.

When $f : M \rightarrow \mathbb{R}$ is a Morse-Bott function there is no known way to express these differentials in terms of the moduli spaces of gradient flow lines of $f : M \rightarrow \mathbb{R}$.

Instanton homology and gauge theory

Theorem. (Fukaya 1996) Let N be a connected sum of two homology 3-spheres and $R(N)$ the space of conjugacy classes of $SU(2)$ representations of $\pi_1(N)$. Then $R(N)$ is divided into $R_i(N)$ with $i \in \mathbb{Z}$, and there is a **spectral sequence** with $E_{ij}^1 \cong H_j(R_i; \mathbb{Z})$ such that $E_{ij}^* \implies I_{i+j}(N)$.

Instanton homology and gauge theory

Theorem. (Fukaya 1996) Let N be a connected sum of two homology 3-spheres and $R(N)$ the space of conjugacy classes of $SU(2)$ representations of $\pi_1(N)$. Then $R(N)$ is divided into $R_i(N)$ with $i \in \mathbb{Z}$, and there is a **spectral sequence** with $E_{ij}^1 \cong H_j(R_i; \mathbb{Z})$ such that $E_{ij}^* \implies I_{i+j}(N)$.

Theorem. (Austin-Braam 1996) Suppose N is a 3-manifold such that the Chern-Simons functional may be perturbed to a Morse-Bott function with only reducible critical orbits. With some additional assumptions, certain Donaldson polynomials on a 4-manifold $X = X_1 \cup_N X_2$ vanish.

Symplectic Floer homology and quantum cohomology

Proposition. (Ruan-Tian 1995) Let (M, ω) be a semi-positive symplectic manifold and H be a self-indexing Bott-type Hamiltonian. Then there exists a **spectral sequence** $E_{i,j}^*$ on the upper half plane such that $E_{i,j}^* \implies HF^{i+j}(M, H)$.

Symplectic Floer homology and quantum cohomology

Proposition. (Ruan-Tian 1995) Let (M, ω) be a semi-positive symplectic manifold and H be a self-indexing Bott-type Hamiltonian. Then there exists a **spectral sequence** $E_{i,j}^*$ on the upper half plane such that $E_{i,j}^* \implies HF^{i+j}(M, H)$.

This includes the cases where H is nondegenerate and where $H = 0$.

Symplectic Floer homology and quantum cohomology

Proposition. (Ruan-Tian 1995) Let (M, ω) be a semi-positive symplectic manifold and H be a self-indexing Bott-type Hamiltonian. Then there exists a **spectral sequence** $E_{i,j}^*$ on the upper half plane such that $E_{i,j}^* \implies HF^{i+j}(M, H)$.

This includes the cases where H is nondegenerate and where $H = 0$.

Theorem. (Liu-Tian 1999) For any compact symplectic manifold, Floer homology equipped with either the intrinsic or exterior product is isomorphic to quantum homology equipped with the quantum product as a ring.

Cascades (Frauenfelder 2004 - Salamon ?)

Let $f : M \rightarrow \mathbb{R}$ be a Morse-Bott function and suppose

$$\text{Cr}(f) = \coprod_{j=1}^l C_j,$$


where C_1, \dots, C_l are disjoint connected critical submanifolds of Morse-Bott index $\lambda_1, \dots, \lambda_l$ respectively. Let $f_j : C_j \rightarrow \mathbb{R}$ be a Morse function on the critical submanifold C_j for all $j = 1, \dots, l$.

Definition

If $q \in C_j$ is a critical point of the Morse function $f_j : C_j \rightarrow \mathbb{R}$ for some $j = 1, \dots, l$, then the **total index** of q , denoted λ_q , is defined to be the sum of the Morse-Bott index of C_j and the Morse index of q relative to f_j , i.e.

$$\lambda_q = \lambda_j + \lambda_q^j.$$

A 3-cascade



For $q \in \text{Cr}(f_j)$, $p \in \text{Cr}(f_i)$, and $n \in \mathbb{N}$, a **flow line with n cascades from q to p** is a $2n - 1$ -tuple:

$$((x_k)_{1 \leq k \leq n}, (t_k)_{1 \leq k \leq n-1})$$

where $x_k \in C^\infty(\mathbb{R}, M)$ and $t_k \in \mathbb{R}_+ = \{t \in \mathbb{R} \mid t \geq 0\}$ satisfy the following for all k .

For $q \in \text{Cr}(f_j)$, $p \in \text{Cr}(f_i)$, and $n \in \mathbb{N}$, a **flow line with n cascades from q to p** is a $2n - 1$ -tuple:

$$((x_k)_{1 \leq k \leq n}, (t_k)_{1 \leq k \leq n-1})$$

where $x_k \in C^\infty(\mathbb{R}, M)$ and $t_k \in \mathbb{R}_+ = \{t \in \mathbb{R} \mid t \geq 0\}$ satisfy the following for all k .

1. Each x_k is a non-constant gradient flow line of f , i.e.

$$\frac{d}{dt}x_k(t) = -(\nabla f)(x_k(t)).$$

For $q \in \text{Cr}(f_j)$, $p \in \text{Cr}(f_i)$, and $n \in \mathbb{N}$, a **flow line with n cascades from q to p** is a $2n - 1$ -tuple:

$$((x_k)_{1 \leq k \leq n}, (t_k)_{1 \leq k \leq n-1})$$

where $x_k \in C^\infty(\mathbb{R}, M)$ and $t_k \in \mathbb{R}_+ = \{t \in \mathbb{R} \mid t \geq 0\}$ satisfy the following for all k .

1. Each x_k is a non-constant gradient flow line of f , i.e.

$$\frac{d}{dt}x_k(t) = -(\nabla f)(x_k(t)).$$

2. For the first cascade $x_1(t)$ we have

$$\lim_{t \rightarrow -\infty} x_1(t) \in W_{f_j}^u(q) \subseteq C_j,$$

and for the last cascade $x_n(t)$ we have

$$\lim_{t \rightarrow \infty} x_n(t) \in W_{f_i}^s(p) \subseteq C_i.$$

3. For $1 \leq k \leq n - 1$ there are critical submanifolds C_{j_k} and gradient flow lines $y_k \in C^\infty(\mathbb{R}, C_{j_k})$ of f_{j_k} , i.e.

$$\frac{d}{dt}y_k(t) = -(\nabla f_{j_k})(y_k(t)),$$

such that $\lim_{t \rightarrow \infty} x_k(t) = y_k(0)$ and
 $\lim_{t \rightarrow -\infty} x_{k+1}(t) = y_k(t_k)$.

3. For $1 \leq k \leq n - 1$ there are critical submanifolds C_{j_k} and gradient flow lines $y_k \in C^\infty(\mathbb{R}, C_{j_k})$ of f_{j_k} , i.e.

$$\frac{d}{dt}y_k(t) = -(\nabla f_{j_k})(y_k(t)),$$

such that $\lim_{t \rightarrow \infty} x_k(t) = y_k(0)$ and
 $\lim_{t \rightarrow -\infty} x_{k+1}(t) = y_k(t_k)$.

Definition

Denote the space of flow lines from q to p with n cascades by $W_n^c(q, p)$, and denote the quotient of $W_n^c(q, p)$ by the action of \mathbb{R}^n by $\mathcal{M}_n^c(q, p) = W_n^c(q, p)/\mathbb{R}^n$. The **set of unparameterized flow lines with cascades from q to p** is defined to be

$$\mathcal{M}^c(q, p) = \bigcup_{n \in \mathbb{Z}_+} \mathcal{M}_n^c(q, p)$$

where $\mathcal{M}_0^c(q, p) = W_0^c(q, p)/\mathbb{R}$.

The \mathbb{Z}_2 -cascade chain complex

Define the k^{th} chain group $C_k^c(f)$ to be the free abelian group generated by the critical points of total index k of the Morse-Smale functions f_j for all $j = 1, \dots, l$, and define $n^c(q, p; \mathbb{Z}_2)$ to be the number of flow lines with cascades between a critical point q of total index k and a critical point p of total index $k - 1$ counted mod 2. Let

$$C_*^c(f) \otimes \mathbb{Z}_2 = \bigoplus_{k=0}^m C_k^c(f) \otimes \mathbb{Z}_2$$

and define a homomorphism $\partial_k^c : C_k^c(f) \otimes \mathbb{Z}_2 \rightarrow C_{k-1}^c(f) \otimes \mathbb{Z}_2$ by

$$\partial_k^c(q) = \sum_{p \in Cr(f_{k-1})} n^c(q, p; \mathbb{Z}_2) p.$$

The pair $(C_*^c(f) \otimes \mathbb{Z}_2, \partial_*^c)$ is called the **cascade chain complex** with \mathbb{Z}_2 coefficients.

The Arnold-Givental conjecture

Let (M, ω) be a $2n$ -dimensional compact symplectic manifold, $L \subset M$ a compact Lagrangian submanifold, and $R \in \text{Diff}(M)$ an antisymplectic involution, i.e. $R^*\omega = -\omega$ and $R^2 = \text{id}$, whose fixed point set is L .

Conjecture. Let H_t be a smooth family of Hamiltonian functions on M for $0 \leq t \leq 1$ and denote by Φ_H the time-1 map of the flow of the Hamiltonian vector field of H_t . If L intersects $\Phi_H(L)$ transversally, then

$$\#(L \cap \Phi_H(L)) \geq \sum_{k=0}^n b_k(L; \mathbb{Z}_2).$$

Proved by Frauenfelder for a class of Lagrangians in Marsden-Weinstein quotients by letting $H \rightarrow 0$ (2004).

The Yang-Mills gradient flow

Let (Σ, g) be a closed oriented Riemann surface, G a compact Lie group, \mathfrak{g} its Lie algebra, and P a principal G -bundle over Σ . Pick an ad-invariant inner product on \mathfrak{g} , let $\mathcal{A}(P)$ denote the affine space of \mathfrak{g} -valued connection 1-forms on P , and define $\mathcal{YM} : \mathcal{A}(P) \rightarrow \mathbb{R}$ by

$$\mathcal{YM}(A) = \int_{\Sigma} F_A \wedge *F_A$$

where $F_A = dA + \frac{1}{2}[A \wedge A]$ is the curvature of A .

The Yang-Mills function is a Morse-Bott function studied by Atiyah-Bott and by Swoboda (2011) using cascades.

Closed Reeb orbits

Let M be a compact, orientable manifold of dimension $2n - 1$ with contact form α . The **Reeb vector field** R_α associated to the contact form α is characterized by

$$\begin{aligned}d\alpha(R_\alpha, -) &= 0 \\ \alpha(R_\alpha) &= 1.\end{aligned}$$

Closed trajectories of the Reeb vector field are critical points of the action functional $\mathcal{A} : C^\infty(S^1, M) \rightarrow \mathbb{R}$

$$\mathcal{A}(\gamma) = \int_\gamma \alpha.$$

Lemma. For any contact structure ξ on M , there exists a contact form α for ξ such that all closed orbits of R_α are nondegenerate.

Contact homology

Let \mathbf{A} be the graded supercommutative algebra freely generated by the “good” closed Reeb orbits over the graded ring $\mathbb{Q}[H_2(M; \mathbb{Z})/\mathcal{R}]$, i.e. $\gamma_1\gamma_2 = (-1)^{|\gamma_1||\gamma_2|}\gamma_2\gamma_1$.

Theorem. (Eliashberg-Hofer 2000) There is a differential $d : \mathbf{A} \rightarrow \mathbf{A}$ defined by counting J -holomorphic curves in the symplectization $(\mathbb{R} \times M, d(e^t\alpha))$ such that (\mathbf{A}, d) is a **differential graded algebra**. Moreover, $HC_*(M, \xi) \stackrel{\text{def}}{=} H_*(\mathbf{A}, d)$ is an invariant of the contact structure ξ .

Theorem. (Bourgeois 2002) Assume that α is a contact form of **Morse-Bott type** for (M, ξ) and that J is an almost complex structure on the symplectization that is S^1 -invariant along the critical submanifolds N_T . Then there is a chain complex with a boundary operator defined by counting **cascades** whose homology is isomorphic to the contact homology $HC_*(M, \xi)$.

Viterbo's symplectic homology

Definition

A compact symplectic manifold (W, ω) has **contact type** boundary if and only if there exists a vector field X defined in a neighborhood of $M = \partial W$ transverse and pointing outward along M such that $\mathcal{L}_X \omega = \omega$.

In this case, $\lambda = \omega(X, \cdot)|_M$ is a contact form on M , and the symplectic homology of W combines the 1-periodic orbits of a Hamiltonian on W with the Reeb orbits on $M = \partial W$.

Bourgeois and Oancea have defined the cascade chain complex for a time-independent Hamiltonian on W whose 1-periodic orbits are transversally nondegenerate (2009). They have also proved that there is an exact sequence relating the symplectic homology groups of W with the linearized contact homology groups of M (2009).

The explicit perturbation of $f : M \rightarrow \mathbb{R}$

Let T_j be a small tubular neighborhood around each connected component $C_j \subseteq Cr(f)$ for all $j = 1, \dots, l$. Pick a positive Morse function $f_j : C_j \rightarrow \mathbb{R}$ and extend f_j to a function on T_j by making f_j constant in the direction normal to C_j for all $j = 1, \dots, l$.

Let $\tilde{T}_j \subset T_j$ be a smaller tubular neighborhood of C_j with the same coordinates as T_j , and let ρ_j be a smooth bump function which is constant in the coordinates parallel to C_j , equal to 1 on \tilde{T}_j , equal to 0 outside of T_j , and decreases on $T_j - \tilde{T}_j$ as the coordinates move away from C_j . For small $\varepsilon > 0$ (and a careful choice of the metric) this determines a Morse-Smale function

$$h_\varepsilon = f + \varepsilon \left(\sum_{j=1}^l \rho_j f_j \right).$$

Identical chain groups

For every sufficiently small $\varepsilon > 0$ and $k = 0, \dots, m$ we have

$$C_k^c(f) \approx C_k(h_\varepsilon) = \bigoplus_{\lambda_j + n = k} C_n(f_j).$$

Identical chain groups

For every sufficiently small $\varepsilon > 0$ and $k = 0, \dots, m$ we have

$$C_k^c(f) \approx C_k(h_\varepsilon) = \bigoplus_{\lambda_j + n = k} C_n(f_j).$$

Is $\mathcal{M}^c(q, p) \approx \mathcal{M}_{h_\varepsilon}(q, p)$ when $\lambda_q - \lambda_p = 1$?

Identical chain groups

For every sufficiently small $\varepsilon > 0$ and $k = 0, \dots, m$ we have

$$C_k^c(f) \approx C_k(h_\varepsilon) = \bigoplus_{\lambda_j + n = k} C_n(f_j).$$

Is $\mathcal{M}^c(q, p) \approx \mathcal{M}_{h_\varepsilon}(q, p)$ when $\lambda_q - \lambda_p = 1$?

If so, then we can use the orientations on $\mathcal{M}_{h_\varepsilon}(q, p)$ to define the cascade chain complex over \mathbb{Z} so that $\partial_k^c = -\partial_k$ for all $k = 0, \dots, m$, where ∂_k is the Morse-Smale-Witten boundary operator of h_ε .

Identical chain groups

For every sufficiently small $\varepsilon > 0$ and $k = 0, \dots, m$ we have

$$C_k^c(f) \approx C_k(h_\varepsilon) = \bigoplus_{\lambda_j + n = k} C_n(f_j).$$

Is $\mathcal{M}^c(q, p) \approx \mathcal{M}_{h_\varepsilon}(q, p)$ when $\lambda_q - \lambda_p = 1$?

If so, then we can use the orientations on $\mathcal{M}_{h_\varepsilon}(q, p)$ to define the cascade chain complex over \mathbb{Z} so that $\partial_k^c = -\partial_k$ for all $k = 0, \dots, m$, where ∂_k is the Morse-Smale-Witten boundary operator of h_ε . In particular,

$$H_*((C_*(f), \partial_*^c)) \approx H_*(M; \mathbb{Z}).$$

Theorem (Banyaga-H 2011)

Assume that f satisfies the Morse-Bott-Smale transversality condition with respect to the Riemannian metric g on M , $f_k : C_k \rightarrow \mathbb{R}$ satisfies the Morse-Smale transversality condition with respect to the restriction of g to C_k for all $k = 1, \dots, l$, and the unstable and stable manifolds $W_{f_j}^u(q)$ and $W_{f_i}^s(p)$ are transverse to the beginning and endpoint maps.

1. When $n = 0, 1$ the set $\mathcal{M}_n^c(q, p)$ is either empty or a smooth manifold without boundary.
2. For $n > 1$ the set $\mathcal{M}_n^c(q, p)$ is either empty or a smooth manifold with corners.
3. The set $\mathcal{M}^c(q, p)$ is either empty or a smooth manifold without boundary.

In each case the dimension of the manifold is $\lambda_q - \lambda_p - 1$. The above manifolds are orientable when M and C_k are orientable.

Compactness

Denote the space of nonempty closed subsets of $M \times \overline{\mathbb{R}}^l$ in the topology determined by the Hausdorff metric by $\mathcal{P}^c(M \times \overline{\mathbb{R}}^l)$, and map a broken flow line with cascades (v_1, \dots, v_n) to its image $\text{Im}(v_1, \dots, v_n) \subset M$ and the time t_j spent flowing along or resting on each critical submanifold C_j for all $j = 1, \dots, l$.

Theorem (Banyaga-H 2011)

The space $\overline{\mathcal{M}}^c(q, p)$ of broken flow lines with cascades from q to p is compact, and there is a continuous embedding

$$\mathcal{M}^c(q, p) \hookrightarrow \overline{\mathcal{M}}^c(q, p) \subset \mathcal{P}^c(M \times \overline{\mathbb{R}}^l).$$

Hence, every sequence of unparameterized flow lines with cascades from q to p has a subsequence that converges to a broken flow line with cascades from q to p .

Correspondence of moduli spaces

Theorem (Banyaga-H 2011)

Let $p, q \in Cr(h_\varepsilon)$ with $\lambda_q - \lambda_p = 1$. For any sufficiently small $\varepsilon > 0$ there is a bijection between unparameterized cascades and unparameterized gradient flow lines of the Morse-Smale function $h_\varepsilon : M \rightarrow \mathbb{R}$ between q and p ,

$$\mathcal{M}^c(q, p) \leftrightarrow \mathcal{M}_{h_\varepsilon}(q, p).$$

Correspondence of moduli spaces

Theorem (Banyaga-H 2011)


Let $p, q \in \text{Cr}(h_\varepsilon)$ with $\lambda_q - \lambda_p = 1$. For any sufficiently small $\varepsilon > 0$ there is a bijection between unparameterized cascades and unparameterized gradient flow lines of the Morse-Smale function $h_\varepsilon : M \rightarrow \mathbb{R}$ between q and p ,

$$\mathcal{M}^c(q, p) \leftrightarrow \mathcal{M}_{h_\varepsilon}(q, p).$$

Definition

Let $p, q \in \text{Cr}(h_\varepsilon)$ with $\lambda_q - \lambda_p = 1$, define an orientation on the zero dimensional manifold $\mathcal{M}^c(q, p)$ by identifying it with the left hand boundary of $\mathcal{M}_{h_\varepsilon}(q, p) \times [0, \varepsilon]$.

Main idea: The Exchange Lemma



The Morse-Bott chain complex with cascades

Define the k^{th} chain group $C_k^c(f)$ to be the free abelian group generated by the critical points of total index k of the Morse-Smale functions f_j for all $j = 1, \dots, l$, and define $n^c(q, p)$ to be the number of flow lines with cascades between a critical point q of total index k and a critical point p of total index $k - 1$ counted with signs determined by the orientations. Let

$$C_*^c(f) = \bigoplus_{k=0}^m C_k^c(f)$$

and define a homomorphism $\partial_k^c : C_k^c(f) \rightarrow C_{k-1}^c(f)$ by

$$\partial_k^c(q) = \sum_{p \in Cr(f_{k-1})} n^c(q, p)p.$$

Correspondence of chain complexes

Theorem (Banyaga-H 2011)

For $\varepsilon > 0$ sufficiently small we have $C_k^c(f) = C_k(h_\varepsilon)$ and $\partial_k^c = -\partial_k$ for all $k = 0, \dots, m$, where ∂_k denotes the Morse-Smale-Witten boundary operator determined by the Morse-Smale function h_ε . In particular, $(C_^c(f), \partial_*^c)$ is a chain complex whose homology is isomorphic to the singular homology $H_*(M; \mathbb{Z})$.*

Correspondence of chain complexes

Theorem (Banyaga-H 2011)

For $\varepsilon > 0$ sufficiently small we have $C_k^c(f) = C_k(h_\varepsilon)$ and $\partial_k^c = -\partial_k$ for all $k = 0, \dots, m$, where ∂_k denotes the Morse-Smale-Witten boundary operator determined by the Morse-Smale function h_ε . In particular, $(C_^c(f), \partial_*^c)$ is a chain complex whose homology is isomorphic to the singular homology $H_*(M; \mathbb{Z})$.*

Moral: The cascade chain complex of a Morse-Bott function $f : M \rightarrow \mathbb{R}$ is **the same** as the Morse-Smale-Witten complex of a small perturbation of f .

Multicomplexes

Let R be a principal ideal domain. A first quadrant **multicomplex** X is a bigraded R -module $\{X_{p,q}\}_{p,q \in \mathbb{Z}_+}$ with differentials

$$d_i : X_{p,q} \rightarrow X_{p-i,q+i-1} \quad \text{for all } i = 0, 1, \dots$$


that satisfy

$$\sum_{i+j=n} d_i d_j = 0 \quad \text{for all } n.$$

A first quadrant multicomplex can be **assembled** to form a filtered chain complex $((CX)_*, \partial)$ by summing along the diagonals, i.e.

$$(CX)_n \equiv \bigoplus_{p+q=n} X_{p,q} \quad \text{and} \quad F_s(CX)_n \equiv \bigoplus_{\substack{p+q=n \\ p \leq s}} X_{p,q}$$

and $\partial_n = d_0 \oplus \dots \oplus d_n$ for all $n \in \mathbb{Z}_+$. The above relations then imply that $\partial_n \circ \partial_{n+1} = 0$ and $\partial_n(F_s(CX)_*) \subseteq F_s(CX)_*$.



A bicomplex has two filtrations, but a general multicomplex only has one filtration.

The bigraded module associated to the filtration

$$F_s(CX)_n \equiv \bigoplus_{\substack{p+q=n \\ p \leq s}} X_{p,q}$$

is

$$G((CX)_*)_{s,t} = F_s(CX)_{s+t} / F_{s-1}(CX)_{s+t} \approx X_{s,t}$$

for all $s, t \in \mathbb{Z}_+$, and the E^1 term of the associated spectral sequence is given by

$$E_{s,t}^1 = Z_{s,t}^1 / (Z_{s-1,t+1}^0 + \partial Z_{s,t+1}^0)$$

where

$$Z_{s,t}^1 = \{c \in F_s(CX)_{s+t} \mid \partial c \in F_{s-1}(CX)_{s+t-1}\}$$

$$Z_{s,t}^0 = \{c \in F_s(CX)_{s+t} \mid \partial c \in F_s(CX)_{s+t-1}\} = F_s(CX)_{s+t}.$$

$E_{s,t}^1$ and d^1 are induced from d_0 and d_1

Theorem

Let $(\{X_{p,q}\}_{p,q \in \mathbb{Z}_+}, \{d_i\}_{i \in \mathbb{Z}_+})$ be a first quadrant multicomplex and $((CX)_*, \partial)$ the associated assembled chain complex. Then the E^1 term of the spectral sequence associated to the filtration of $(CX)_*$ determined by the restriction $p \leq s$ is given by $E_{s,t}^1 \approx H_{s+t}(X_{s,*}, d_0)$ where $(X_{s,*}, d_0)$ denotes the following chain complex.

$$\dots \xrightarrow{d_0} X_{s,3} \xrightarrow{d_0} X_{s,2} \xrightarrow{d_0} X_{s,1} \xrightarrow{d_0} X_{s,0} \xrightarrow{d_0} 0$$

Moreover, the d^1 differential on the E^1 term of the spectral sequence is induced from the homomorphism d_1 in the multicomplex.

However, d^r is not induced from d_r for $r \geq 2$

Consider the following first quadrant double complex

$$\begin{array}{ccccc}
 0 & \xleftarrow{d_1} & 0 & \xleftarrow{d_1} & 0 \\
 \downarrow d_0 & & \downarrow d_0 & & \downarrow d_0 \\
 \langle x_{0,1} \rangle & \xleftarrow{d_1} & \langle x_{1,1} \rangle & \xleftarrow{d_1} & 0 \\
 \downarrow d_0 & & \downarrow d_0 & & \downarrow d_0 \\
 0 & \xleftarrow{d_1} & \langle x_{1,0} \rangle & \xleftarrow{d_1} & \langle x_{2,0} \rangle
 \end{array}$$

where $\langle x_{p,q} \rangle$ denotes the free abelian group generated by $x_{p,q}$, the groups $X_{p,q} = 0$ for $p + q > 2$, and the homomorphisms d_0 and d_1 satisfy the following: $d_0(x_{1,1}) = x_{1,0}$, $d_1(x_{1,1}) = x_{0,1}$, and $d_1(x_{2,0}) = x_{1,0}$. In this case, $d_2 = 0$ but $d^2 \neq 0$

The Morse-Bott-Smale multicomplex

Let $C_p(B_i)$ be the group of “ p -dimensional chains” in the critical submanifolds of index i . Assume that $f : M \rightarrow \mathbb{R}$ is a Morse-Bott-Smale function and the manifold M , the critical submanifolds, and their negative normal bundles are all orientable.

If $\sigma : P \rightarrow B_i$ is a singular C_p -space in $S_p^\infty(B_i)$, then for any $j = 1, \dots, i$ composing the projection map π_2 onto the second component of $P \times_{B_i} \overline{\mathcal{M}}(B_i, B_{i-j})$ with the endpoint map $\partial_+ : \overline{\mathcal{M}}(B_i, B_{i-j}) \rightarrow B_{i-j}$ gives a map

$$P \times_{B_i} \overline{\mathcal{M}}(B_i, B_{i-j}) \xrightarrow{\pi_2} \overline{\mathcal{M}}(B_i, B_{i-j}) \xrightarrow{\partial_+} B_{i-j}.$$

$$\begin{array}{ccccccc}
 \dots & & \vdots & & & & \\
 & & \oplus & & & & \\
 \dots & C_1(B_2) & \xrightarrow{\partial_0} & C_0(B_2) & \xrightarrow{\partial_0} & 0 & \\
 & \searrow^{\partial_1} & & \searrow^{\partial_1} & & & \\
 & \oplus & & \oplus & & \oplus & \\
 \dots & C_2(B_1) & \xrightarrow{\partial_0} & C_1(B_1) & \xrightarrow{\partial_0} & C_0(B_1) & \xrightarrow{\partial_0} & 0 \\
 & \searrow^{\partial_1} & & \searrow^{\partial_1} & & \searrow^{\partial_1} & & \\
 & \oplus & & \oplus & & \oplus & & \oplus \\
 \dots & C_3(B_0) & \xrightarrow{\partial_0} & C_2(B_0) & \xrightarrow{\partial_0} & C_1(B_0) & \xrightarrow{\partial_0} & C_0(B_0) & \xrightarrow{\partial_0} & 0 \\
 & \parallel & & \parallel & & \parallel & & \parallel & & \\
 \dots & C_3(f) & \xrightarrow{\partial} & C_2(f) & \xrightarrow{\partial} & C_1(f) & \xrightarrow{\partial} & C_0(f) & \xrightarrow{\partial} & 0
 \end{array}$$

The Morse-Bott Homology Theorem

Theorem (Banyaga-H 2010)

The homology of the Morse-Bott-Smale multicomplex $(C_(f), \partial)$ is independent of the Morse-Bott-Smale function $f : M \rightarrow \mathbb{R}$.*

Therefore,

$$H_*(C_*(f), \partial) \approx H_*(M; \mathbb{Z}).$$

Note: If f is constant, then $(C_*(f), \partial)$ is the chain complex of singular N -cube chains. If f is Morse-Smale, then $(C_*(f), \partial)$ is the Morse-Smale-Witten chain complex. This gives a new proof of the Morse Homology Theorem.

References

- ▶ David Austin and Peter Braam, *Morse-Bott theory and equivariant cohomology*, **The Floer memorial volume**, Progr. Math. **133** (1995), 123–183.
- ▶ David Austin and Peter Braam, *Equivariant Floer theory and gluing Donaldson polynomials*, Topology **35** (1996), no. 1, 167–200.
- ▶ Augustin Banyaga and David Hurtubise, *A proof of the Morse-Bott Lemma*, Expo. Math. **22**, no. 4, 365–373, 2004.
- ▶ Augustin Banyaga and David Hurtubise, **Lectures on Morse homology**, Kluwer Texts in the Mathematical Sciences **29**, 2004.
- ▶ Augustin Banyaga and David Hurtubise, *The Morse-Bott inequalities via a dynamical systems approach*, Ergodic Theory Dynam. Systems **29** (2009), no. 6, 1693–1703.

References

- ▶ Augustin Banyaga and David Hurtubise, *Morse-Bott homology*, Trans. Amer. Math. Soc., **362** (2010), no. 8, 3997–4043.
- ▶ Augustin Banyaga and David Hurtubise, *Cascades and perturbed Morse-Bott functions*, preprint, 2011.
- ▶ Frédéric Bourgeois and Alexandru Oancea, *Symplectic homology, autonomous Hamiltonians, and Morse-Bott moduli spaces*, Duke Math. J. **146** (2009), no. 1, 74–174.
- ▶ Frédéric Bourgeois and Alexandru Oancea, *An exact sequence for contact and symplectic homology*, Invent. Math. **179** (2009), no. 3, 611–680.
- ▶ Andreas Floer, *An instanton-invariant for 3-manifolds*, Comm. Math. Phys. **118** (1988), no. 2, 215–240.

References

- ▶ Andreas Floer, *Instanton homology, surgery, and knots*, **Geometry of low-dimensional manifolds**, London Math. Soc. Lecture Note Ser. **150** (1990), 97–114.
- ▶ Andreas Floer, *Morse theory for Lagrangian intersections*, *Journal of Differential Geom* **28** (1988), 513–547.
- ▶ Urs Frauenfelder, *The Arnold-Givental conjecture and moment Floer homology*, *Int. Math. Res. Not.* **2004**, no. 42, 2179–2269.
- ▶ Kenji Fukaya, *Floer homology of connected sum of homology 3-spheres*, *Topology* **35** (1996), no. 1, 89–136.
- ▶ David Hurtubise, *Multicomplexes and spectral sequences*, *J. Algebra Appl.* **9** (2010), no. 4, 519–530.

References

- ▶ Jean-Pierre Meyer, *Acyclic models for multicomplexes*, Duke Math. J. **45** (1978), no. 1, 67–85.
- ▶ Yongbin Ruan and Gang Tian, *Bott-type symplectic Floer cohomology and its multiplication structures*, Math. Res. Lett. **2** (1995), no. 2, 203–219.
- ▶ Jan Swoboda, *Morse homology for the Yang-Mills gradient flow*, arXiv:1103.0845v1, 2011.
- ▶ Gang Liu and Gang Tian, *On the equivalence of multiplicative structures in Floer homology and quantum homology*, Acta Math. Sin. **15** (1999), no. 1, 53–80.
- ▶ Joa Weber, *The Morse-Witten complex via dynamical systems*, Expo. Math. **24** (2006), no. 2, 127–159.
- ▶ Edward Witten. **Supersymmetry and Morse theory**, J. Differential Geom. **17** (1982), no. 4, 661–692 (1983).